

For adults and children 12 years and older who weigh at least 88 pounds with moderate to severe plaque psoriasis

PATIENT BROCHURE

 **ICOTYDE™**
(icotrokinra) tablets



NOW APPROVED

A new kind of treatment in a once-daily pill


Pill not actual size.

ICOTYDE™ 200 mg is a prescription medicine used to treat moderate to severe plaque psoriasis in adults and children 12 years of age and older who weigh at least 88 pounds (40 kg), who may benefit from taking injections or medicines by mouth (systemic therapy) or treatment using ultraviolet or UV light (phototherapy).

SELECTED IMPORTANT SAFETY INFORMATION

ICOTYDE™ is not for everyone; only your healthcare provider can decide if it's right for you. It is a prescription medicine that may cause serious side effects, including infections. It may increase the risk of infections and lower your ability to fight them. Tell your healthcare provider if you have an infection, tuberculosis (TB), or if you need a vaccine. Please read the Important Safety Information on [pages 8-9](#) and [Medication Guide](#) to learn more about these and other risks for ICOTYDE. Discuss any questions you have with your healthcare provider.

THE FIRST & ONLY IL-23R TARGETED ORAL PEPTIDE IN A ONCE-DAILY PILL



In plaque psoriasis, interleukin-23 (IL-23) is a protein that plays a key role in inflammation.

When IL-23 binds with its receptor, inflammation can occur in the body.

ICOTYDE targets the IL-23 receptor which **blocks** IL-23 from **binding**. This **prevents** the release of inflammatory proteins.*

*The clinical significance of these findings is unknown.

PROVEN SAFETY PROFILE

ICOTYDE has been extensively studied in people 12 and older

Consistent safety profile across

2300+
people
in

4 clinical trials

SELECTED IMPORTANT SAFETY INFORMATION

ICOTYDE™ is not for everyone; only your healthcare provider can decide if it's right for you. It is a prescription medicine that may cause serious side effects, including infections. It may increase the risk of infections and lower your ability to fight them. Tell your healthcare provider if you have an infection, tuberculosis (TB), or if you need a vaccine. Please read the Important Safety Information on [pages 8-9](#) and [Medication Guide](#) to learn more about these and other risks for ICOTYDE. Discuss any questions you have with your healthcare provider.

Most common side effects at 4 months (16 weeks)*

	ICOTYDE	Placebo†
Headache	4.1%	3.3%
Nausea	1.2%	0.5%
Cough	1.2%	0.2%
Fungal infection		
Tinea	0.5%	0.0%
Other‡	0.8%	0.0%
Tiredness	1.0%	0.5%

These are not all the possible side effects of ICOTYDE. Talk to your doctor about all possible side effects.

*In 4 clinical studies, 1296 people took ICOTYDE, and 568 people took a placebo.

†A placebo is a pill with no medicine in it but looks the same and is taken the same way as the medicine being tested.

‡Includes oral candidiasis, onychomycosis, skin candida, urinary tract candidiasis, vulvovaginal candidiasis, fungal skin infection, genital infection fungal, ear infection fungal, and laryngitis fungal.

POWERFUL SKIN CLEARANCE



In 2 clinical trials, **more than 5 out of 10 people** saw at least **90% clearer skin** at 4 months (16 weeks). **Results may vary.**

2x as many people had completely clear skin with ICOTYDE compared to another psoriasis pill

At 6 months (24 weeks), over 3 out of 10 people taking ICOTYDE had completely clear skin, compared to less than 2 out of 10 people taking Sotyktu® (deucravacitinib). **Results may vary.**

Sotyktu is a registered trademark of Bristol-Myers Squibb Company.

SELECTED IMPORTANT SAFETY INFORMATION

ICOTYDE™ is not for everyone; only your healthcare provider can decide if it's right for you. It is a prescription medicine that may cause serious side effects, including infections. It may increase the risk of infections and lower your ability to fight them. Tell your healthcare provider if you have an infection, tuberculosis (TB), or if you need a vaccine. Please read the Important Safety Information on [pages 8-9](#) and [Medication Guide](#) to learn more about these and other risks for ICOTYDE.

Discuss any questions you have with your healthcare provider.

CLEARANCE OVER TIME

Percentage of people on ICOTYDE who saw at least 90% clearer skin



Real patients, real results



0 months on ICOTYDE



4 months (16 weeks) on ICOTYDE



6 months (24 weeks) on ICOTYDE

Photos are of a real patient. Images are owned by Janssen Biotech, Inc., a Johnson & Johnson company, from a clinical trial. Individual results may vary.

CLEARER SKIN IN HARD-TO-TREAT AREAS

In a clinical study of people with plaque psoriasis* with at least moderate severity in hard-to-treat areas

Most people (nearly 6 out of 10) saw clear or almost clear skin at 4 months (16 weeks)

After 4 months (16 weeks), patients and healthcare providers knew that ICOTYDE was being used. This may have increased results. **Results may vary.**

SIGNIFICANT ITCH IMPROVEMENT


At 4 months (16 weeks),

~3 out of 5
people

saw
significant **itch**
improvement†

†59% of patients.

Scalp clearance



Genital clearance



*In this study, people had plaque psoriasis on at least 1% of their body surface area.

SELECTED IMPORTANT SAFETY INFORMATION

ICOTYDE™ is not for everyone; only your healthcare provider can decide if it's right for you. It is a prescription medicine that may cause serious side effects, including infections. It may increase the risk of infections and lower your ability to fight them. Tell your healthcare provider if you have an infection, tuberculosis (TB), or if you need a vaccine. Please read the Important Safety Information on [pages 8-9](#) and [Medication Guide](#) to learn more about these and other risks for ICOTYDE. Discuss any questions you have with your healthcare provider.

ONE PILL, ONCE A DAY

Wake + Take

When you wake up, take with water on an empty stomach at least 30 minutes before eating

For full dosing instructions, including how to take ICOTYDE if you have difficulty swallowing pills, please see the [Medication Guide](#).

SELECTED IMPORTANT SAFETY INFORMATION

ICOTYDE™ is not for everyone; only your healthcare provider can decide if it's right for you. It is a prescription medicine that may cause serious side effects, including infections. It may increase the risk of infections and lower your ability to fight them. Tell your healthcare provider if you have an infection, tuberculosis (TB), or if you need a vaccine. Please read the Important Safety Information on [pages 8-9](#) and [Medication Guide](#) to learn more about these and other risks for ICOTYDE. Discuss any questions you have with your healthcare provider.



Pill not actual size.

Once you and your doctor have decided that ICOTYDE is right for you

ICOTYDE *withMe*

Savings Program for eligible commercially insured patients

PAY AS LITTLE AS \$0 PER MONTH*

Here's what to expect



Your doctor sends your prescription to a specialty pharmacy (SP).



Your insurance reviews and confirms your coverage.

This is handled by your insurance company and the SP.



Answer the phone call from the SP to schedule delivery.

Some patients will receive a phone call from "Healthcare" at (888) 796-8154.

Consider saving the SP's number so you'll recognize the call when it comes through.

It's very important you answer the phone call.

*Maximum program benefit per calendar year shall apply. Not valid for patients using Medicare, Medicaid, or other government-funded programs to pay for their medicines. Terms expire at the end of each calendar year. Offer subject to change or end without notice. Restrictions, including monthly maximums, may apply. Participate without sharing your income information. See program requirements at [ICOTYDEwithMeSavings.com](https://www.icotyde.com/withMeSavings).

The support and resources provided by ICOTYDE withMe are not intended to provide medical advice, replace a treatment plan you receive from your doctor or nurse, or serve as a reason for you to start or stay on treatment.



Why do you need a specialty pharmacy?

Medicines for chronic conditions (like plaque psoriasis) often require extra care and support. A specialty pharmacy works closely with you, your doctor, and your insurance company to confirm your access and get you support and then ships the medicine directly to your home.

Once you and your doctor have decided that ICOTYDE is right for you, sign up for

ICOTYDE *withMe*

Get clear, flexible support to help you get started and stay on track with treatment



Dedicated Nurse Guide*

ICOTYDE withMe gives you 1-on-1 support from a Nurse Guide. They're here to answer your questions and connect you to helpful resources.



Prescription and cost support

ICOTYDE withMe can help you verify your insurance coverage, understand how to get your prescriptions filled, and find ways to make treatment more affordable, even if you don't have insurance.



Patient portal

The patient portal lets you view account details and get access to personalized resources all in one place.

Visit [icotyde.com/signup](https://www.icotyde.com/signup) for more information

For any questions or to sign up by phone, call us at **833-withMe1** (833-948-4631), Monday–Friday, 8:00 AM–11:00 PM ET.

The support and resources provided by ICOTYDE withMe are not intended to provide medical advice, replace a treatment plan you receive from your doctor or nurse, or serve as a reason for you to start or stay on treatment.

*Nurse Guides do not provide medical advice. Please ask your doctor any questions you might have about your disease and treatment.

INDICATION

WHAT IS ICOTYDE™ (icotrokinra)?

ICOTYDE™ 200 mg is a prescription medicine used to treat moderate to severe plaque psoriasis in adults and children 12 years of age and older who weigh at least 88 pounds (40 kg), who may benefit from taking injections or medicines by mouth (systemic therapy) or treatment using ultraviolet or UV light (phototherapy).

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about ICOTYDE?

- **Infections.** Medicines that interact with the immune system, such as ICOTYDE, may lower your ability to fight infections and may increase your risk of infections. Your healthcare provider may check you for infections and tuberculosis (TB) before starting treatment and may treat you for TB before you begin treatment with ICOTYDE if you have a history of TB or have active TB. Your healthcare provider should watch you closely for signs and symptoms of TB during and after treatment with ICOTYDE.

Tell your healthcare provider right away if you have any infection or have symptoms of an infection, including:

- fever, sweat, or chills
- cough
- shortness of breath
- blood in your mucus (phlegm)
- muscle aches
- warm, red, or painful skin or sores on your body different from your psoriasis
- weight loss
- diarrhea or stomach pain
- burning when you urinate or urinating more often than normal

Please see additional Important Safety Information on [page 9](#).

IMPORTANT SAFETY INFORMATION (continued)

Before taking ICOTYDE, tell your healthcare provider about all of your medical conditions, including if you:

- have an infection that does not go away or that keeps coming back.
- have tuberculosis (TB) or have been in close contact with someone with TB.
- have recently received or are scheduled to receive an immunization (vaccine). Avoid receiving live vaccines during treatment with ICOTYDE.
- have kidney problems.
- are pregnant or plan to become pregnant. It is not known if ICOTYDE can harm your unborn baby.

Pregnancy Safety Study. There is a pregnancy safety study for women who take ICOTYDE during pregnancy. The purpose of this study is to collect information about the health of you and your baby. If you are pregnant or become pregnant during treatment with ICOTYDE, you can report your pregnancy by calling 1-800-526-7736 or visiting www.ICOTYDE.com.

- are breastfeeding or plan to breastfeed. It is not known if ICOTYDE passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby during treatment with ICOTYDE.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

What are the possible side effects of ICOTYDE?

ICOTYDE may cause serious side effects. See “What is the most important information I should know about ICOTYDE?”

The most common side effects of ICOTYDE include:

- headache
- nausea
- cough
- fungal infection
- tiredness

IMPORTANT SAFETY INFORMATION (continued)

These are not all the possible side effects of ICOTYDE. Call your doctor for medical advice about side effects. **You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.**

How should I take ICOTYDE?

- Take ICOTYDE exactly as your healthcare provider tells you to take it.
- Take ICOTYDE 1 time a day when you wake up on an empty stomach with water. Wait at least 30 minutes after taking ICOTYDE before eating food.
- If you have difficulty swallowing tablets, ICOTYDE can be dispersed in water. For more information, please read the **Medication Guide**.
- If you miss a dose of ICOTYDE, take the dose as soon as you remember and go back to your regular schedule the next day.

Please read the full Prescribing Information, including Medication Guide, for ICOTYDE and discuss any questions that you have with your doctor.

cp-564097v2

For adults and children 12 years and older who weigh at least 88 pounds
with moderate to severe plaque psoriasis

 **ICOTYDE™**
(icotrokinra) tablets

Proven safety profile

Across 2300+ people in 4 clinical trials

Powerful skin clearance

In 2 clinical trials, more than 5 out of 10 people saw
at least 90% clearer skin at 4 months (16 weeks)

One pill, once a day

Visit [icotyde.com](https://www.icotyde.com)

Pill not actual size.



SELECTED IMPORTANT SAFETY INFORMATION

ICOTYDE™ is not for everyone; only your healthcare provider can decide if it's right for you. It is a prescription medicine that may cause serious side effects, including infections. It may increase the risk of infections and lower your ability to fight them. Tell your healthcare provider if you have an infection, tuberculosis (TB), or if you need a vaccine. Please read the Important Safety Information on [pages 8-9](#) and [Medication Guide](#) to learn more about these and other risks for ICOTYDE. Discuss any questions you have with your healthcare provider.